

Supplemental Table S1. Baseline Characteristics of Patients According to the Presence of Distant Metastasis and Persistent/Recurrent Disease

Characteristic	Without distant metastasis (n=94)	Distant metastasis (n=3)	P value	NED (n=89)	Persistent/recurrent disease (n=8)	P value
Age, yr	49.8±13.2	68.5±1.3	<0.01	49.2±13.2	63.3±9.6	<0.01
≥55	33 (35.1)	3 (100)	0.09	30 (33.7)	6 (75.0)	0.05
Male sex	25 (26.6)	1 (33.3)	0.99	23 (25.8)	3 (37.5)	0.77
Primary tumor size, cm	3.0 (1.8–4.3)	2.0 (1.9–5.5)	0.96	3.0 (2.0–4.3)	2.2 (1.6–4.8)	0.82
>4	24 (25.5)	1 (33.3)	0.99	23 (25.8)	2 (25.0)	0.99
Extra-thyroidal extension			<0.01			0.01
Gross	1 (1.1)	2 (66.7)		1 (1.1)	2 (25.0)	
WHO classification			<0.01			<0.01
Minimally invasive	73 (77.7)	0		70 (78.7)	3 (37.5)	
Encapsulated angioinvasive	18 (19.1)	1 (33.3)		17 (19.1)	2 (25.0)	
Widely invasive	3 (3.2)	2 (66.7)		2 (2.2)	3 (37.5)	

Values are expressed as mean ± standard deviation, number (%), or median (interquartile range). NED, no evidence of disease; WHO, World Health Organization.